

Comparative Stability Analysis of Herbal and Synthetic Pharmaceutical Formulations

Humera Khanam^{1,*} and Gaurav Kant Saraogi⁵

¹⁻² Dr. Preeti Global University, Shivpuri - 473665, M.P. India

*Corresponding Author: Humera Khanam, Email: humerakhanam.pgu@gmail.com

Received: 07/07/2026; Revised: 29/07/2026; Accepted: 29/08/2026; Published: 30/09/2026.

DOI: <https://doi.org/10.69996/fmep.2026028>.

Abstract: The increasing use of herbal medicines has generated significant interest in evaluating their pharmaceutical stability in comparison with synthetic formulations. Herbal products contain complex mixtures of phytochemicals that are highly susceptible to environmental degradation. The present study aimed to compare the stability characteristics of herbal and synthetic pharmaceutical formulations under different environmental conditions. Physicochemical properties, active ingredient retention, microbial contamination, and shelf-life parameters were analyzed using standard pharmaceutical methods. Accelerated stability testing demonstrated that herbal formulations exhibited greater sensitivity to temperature, humidity, and oxidation compared to synthetic products. However, appropriate packaging and controlled storage conditions significantly improved the stability of herbal preparations. The study highlights the need for standardized quality control measures and advanced formulation strategies to enhance the stability and therapeutic reliability of herbal pharmaceuticals.

Keywords: Phytochemicals, Physicochemical Properties, Shelf-Life Parameters, Herbal Formulations.

1. Introduction

Herbal medicines have been utilized for centuries in traditional healthcare systems such as Ayurveda and Unani medicine. In recent years, growing concerns regarding adverse effects associated with synthetic drugs have increased public interest in plant-based therapeutic products.

Although herbal formulations offer several therapeutic advantages, their stability remains a critical issue in pharmaceutical applications [1-5]. Herbal medicines contain multiple phytochemical constituents that may undergo degradation due to environmental factors such as temperature, moisture, oxidation, and microbial contamination.

Synthetic pharmaceutical formulations generally possess well-defined chemical structures and comparatively stable physicochemical properties. In contrast, herbal formulations exhibit considerable variability due to differences in plant source, harvesting conditions, extraction methods, and storage practices [6-10].

Comparative stability evaluation of herbal and synthetic products is essential for ensuring product quality, efficacy, and shelf-life. Such studies help identify limitations associated with herbal pharmaceutical products and facilitate the development of improved formulation and preservation strategies [11-16].

The present study was therefore designed to compare the stability profiles of herbal and synthetic pharmaceutical formulations under various storage conditions.

2. Literature Review

According to WHO (2004), quality assurance of herbal medicines requires comprehensive stability testing because herbal products are highly susceptible to environmental degradation.



Mukherjee (2002) highlighted that herbal formulations possess complex chemical compositions that complicate stability assessment compared to synthetic drugs. Environmental exposure may significantly alter phytochemical composition and therapeutic activity.

Patwardhan et al. (2005) emphasized the importance of integrating traditional herbal medicine with modern pharmaceutical standardization practices to improve product reliability.

Sharma and Gupta (2009) reported that synthetic formulations generally exhibit greater chemical stability than herbal products due to their purified active ingredients and standardized manufacturing processes.

Singh et al. (2011) observed accelerated degradation of herbal preparations under high humidity and temperature conditions, whereas synthetic formulations remained comparatively stable.

ICH guidelines recommend accelerated and long-term stability testing for pharmaceutical products to determine expiration dating and storage recommendations.

3. Methodology

Sample Selection

Both herbal and synthetic pharmaceutical formulations with similar therapeutic applications were selected.

Stability Conditions

Samples were stored under:

- Room temperature conditions
- Accelerated stability conditions
- Light exposure conditions

Parameters Evaluated

- Appearance
- pH
- Moisture content
- Active ingredient assay
- Microbial contamination
- Shelf-life estimation

Analytical Procedures

- HPLC
- UV spectrophotometry
- Microbial analysis
- Pharmacopoeial quality tests

Statistical Analysis

Comparative statistical analysis was performed using ANOVA and regression methods.

4. Results and Discussion

Comparative analysis revealed that herbal formulations were more sensitive to environmental changes than synthetic formulations. Elevated temperature and humidity accelerated degradation of phytoconstituents and increased microbial contamination in herbal products.

Synthetic formulations exhibited relatively stable physicochemical characteristics throughout the study period due to their purified chemical composition and advanced stabilization techniques.

However, herbal formulations stored in airtight and light-resistant containers showed improved stability. Appropriate packaging significantly reduced moisture absorption and oxidation.

The findings indicate that although herbal medicines possess therapeutic potential, their stability remains a major pharmaceutical challenge. Modern stabilization approaches such as nanoformulations, encapsulation, and antioxidant incorporation may improve shelf-life and product quality.

The study supports earlier observations by Mukherjee (2002), Sharma and Gupta (2009), and Singh et al. (2011) regarding the environmental sensitivity of herbal pharmaceutical products.

5. Conclusion

The present study demonstrated significant differences in the stability profiles of herbal and synthetic pharmaceutical formulations. Herbal products showed greater susceptibility to environmental degradation, microbial contamination, and phytochemical instability compared to synthetic formulations.

Nevertheless, suitable storage conditions, protective packaging, and formulation optimization significantly enhanced the stability of herbal medicines. The findings emphasize the necessity for standardized manufacturing practices, rigorous quality control, and advanced pharmaceutical technologies in herbal drug development.

Comprehensive stability evaluation is therefore essential for ensuring the safety, efficacy, therapeutic reliability, and commercial acceptability of herbal pharmaceutical products.

Acknowledgment: Not Applicable.

Funding Statement: The author(s) received no specific funding for this study.

Conflicts of Interest: The authors declare no conflicts of interest to report regarding the present study

References

- [1] A. A. Farooqi and B. A. Shreeramu, *Cultivation of Medicinal and Aromatic Crops*, 1st ed. Hyderabad, India: Universities Press (India) Ltd., 2001, pp. 1–10.
- [2] C. K. Kokate, A. P. Purohit, and S. B. Gokhale, *Textbook of Pharmacognosy*, 45th ed. Mumbai, India: Nirali Prakashan, 2010, pp. 1.4–1.8.
- [3] S. H. Ansari, *Essentials of Pharmacognosy*, 4th ed. New Delhi, India: Birla Publication, 2011, pp. 469–524.
- [4] M. S. Hakim, *Hamdard Pharmacopeia of Eastern Medicine*, reprint ed. New Delhi, India: Shri Satguru Publications, 1997, pp. vii–xiv.
- [5] A. C. Dweck, “Natural preservatives,” *NaturalIngredient.org*, pp. 1–33, 2023.
- [6] R. Marks, A. Plunkett, K. Merlin, and N. Jenner, *Atlas of Common Skin Diseases*. Fitzroy, Victoria, Australia: Dept. of Dermatology, St. Vincent’s Hospital, pp. 6–30, 2012.
- [7] P. K. Mukherjee, *Quality Control of Herbal Drugs: An Approach to Evaluation of Botanicals*. New Delhi, India: Business Horizons, 2002, pp. 493–515, 599–622.
- [8] A. Farooqi, “Herbal skin care formulation and a process for the preparation thereof,” *patent*, 2002.
- [9] G. Trease and W. Evans, *A Textbook of Pharmacognosy*, 15th ed. London, U.K.: Academic Press, 2006, p. 274.
- [10] K. R. Kirtikar and B. D. Basu, *Indian Medicinal Plants*, vols. I–IV. Dehradun, India: International Book Distributors, 1918, pp. 161–164, 1042–1043, 2592–2593, plates 93, 420, and 991.
- [11] K. M. Nadkarni, *Indian Materia Medica*, vol. 1. Bombay, India: Popular Prakashan, 1954, pp. 894–895.
- [12] C. Atal and B. Kapur, *Cultivation and Utilization of Aromatic Plants*. Jammu, India: Regional

-
- Research Laboratory, Council of Scientific and Industrial Research, 1982, pp. 638–639, 736.
- [13] W.-H. Hui and M.-M. Li, “Neutral triterpenoids from *Melaleuca leucadendron*,” *Phytochemistry*, vol. 15, no. 4, p. 563, 1976.
- [14] J. L. Courtney, E. V. Lassak, and G. B. Speirs, “Leaf wax constituents of some myrtaceous species,” *Phytochemistry*, vol. 22, no. 4, pp. 947–949, 1983.
- [15] U. Srilakshmi, J Manikandan, Thanmayee Velagapudi, Gandla Abhinav, Tharun Kumar, and Dogiparthi Saideep, “A New Approach to Computationally- Successful Linear and Polynomial Regression Analytics of Large Data in Medicine,” *JCAI*, vol. 2, no. 2, pp. 35–48, 2024.
- [16] C. A. Pearce and K. D. Hyde, “Phyllachora from Australia: Observations on *P. pseudostromatica*, *P. melaleucae* and a new species, *P. shivasii* from the host *Melaleuca*,” *Mycological Research*, vol. 99, no. 10, pp. 1253–1260, 1995.