

Role of Native Pollinators in Enhancing Biodiversity, Crop Productivity, and Ecological Resilience

Rahul Singh Koushal^{1,*} and Popat Prabhakar Pathare⁵

¹⁻²Dr. Preeti Global University, Shivpuri - 473665, M.P. India

*Corresponding Author: Rahul Singh Koushal, Email: rahulsinghkaushal280@gmail.com

Received: 07/07/2026; Revised: 25/07/2026; Accepted: 16/08/2026; Published: 30/09/2026.

DOI: <https://doi.org/10.69996/fmep.2026033>.

Abstract: Native pollinators play a crucial role in maintaining biodiversity, supporting agricultural production, and sustaining ecological resilience. Insects such as bees, butterflies, moths, beetles, and flies contribute significantly to the pollination of wild plants and cultivated crops, thereby ensuring seed production, fruit development, and genetic diversity within ecosystems. Despite their ecological importance, native pollinator populations are declining globally due to habitat loss, pesticide exposure, invasive species, climate change, and intensive agricultural expansion. This study investigates the role of native pollinators in promoting ecosystem sustainability and enhancing crop productivity while assessing the environmental threats that endanger their survival. The paper focuses on the ecological functions of native pollinators and their contributions to food security, ecosystem balance, and plant reproduction. It also examines how pollinator diversity strengthens ecosystem resilience by supporting stable plant-pollinator networks and improving the adaptability of ecosystems to environmental stress. Furthermore, the research explores conservation strategies such as habitat restoration, pollinator-friendly farming, reduction of chemical pesticide use, and the protection of native flowering vegetation. Through an analysis of ecological studies and agricultural evidence, this paper demonstrates that conserving native pollinators is essential for sustaining biodiversity and ensuring long-term agricultural and environmental health. The study concludes that effective pollinator conservation policies are necessary to maintain ecological integrity and support sustainable development in the face of growing environmental challenges.

Keywords: Habitat Loss, Pesticide Exposure, Invasive Species, Climate Change, Pollinator-Friendly Farming.

1. Introduction

Native pollinators are essential components of terrestrial ecosystems and agricultural landscapes. Bees, butterflies, moths, flies, beetles, birds, and bats contribute significantly to the pollination of both cultivated crops and wild plant species. Their interactions with flowering plants support seed formation, fruit production, and genetic diversity, thereby sustaining ecosystem functioning and food production systems. In natural ecosystems, pollinators facilitate plant reproduction, which in turn provides food and habitat resources for numerous animal species [1-5]. In agricultural systems, pollination services increase crop yield, improve fruit quality, and enhance nutritional diversity. Despite their ecological and economic importance, native pollinator populations are declining rapidly due to habitat destruction, climate change, pesticide exposure, invasive species, and intensive farming practices.

Native pollinators are often more efficient and better adapted to local environmental conditions than introduced commercial pollinators. Solitary bees, bumblebees, and indigenous insect species possess specialized behavioral and physiological traits that enable effective pollination of regional crops and native plants. Their contributions are particularly important in maintaining ecological resilience, as diverse pollinator communities provide greater stability



against environmental disturbances and climate variability [6-10]. However, modern agricultural systems frequently prioritize short-term productivity over biodiversity conservation, resulting in habitat simplification and reduced floral diversity. The replacement of heterogeneous landscapes with monocultures has diminished nesting sites and food resources necessary for native pollinator survival.

The decline of native pollinators threatens both biodiversity and ecosystem resilience. Reduced pollinator diversity can lead to lower plant reproductive success, decreased ecosystem productivity, and weakened ecological interactions. Furthermore, the disappearance of pollinators may affect entire food chains by limiting resources for herbivores, seed-eating organisms, and higher trophic levels [11-15]. Conservation of native pollinators is therefore critical for sustaining ecological balance and ensuring long-term agricultural productivity. Strategies such as habitat restoration, creation of pollinator-friendly landscapes, reduction of pesticide use, and promotion of native flowering plants can help restore pollinator populations and strengthen ecosystem sustainability. This research seeks to investigate the ecological role of native pollinators in supporting biodiversity, enhancing crop productivity, and improving ecosystem resilience under changing environmental conditions [16-20].

Native pollinators are fundamental to the maintenance of healthy ecosystems and sustainable agricultural systems. They provide essential ecological services by facilitating the pollination of flowering plants, which supports plant reproduction, seed formation, and fruit production. Pollination is a critical ecological process that enables genetic exchange among plants and contributes to the stability and diversity of natural ecosystems. Native pollinators include a wide range of organisms such as solitary bees, bumblebees, butterflies, moths, beetles, flies, birds, and bats [21-26]. These organisms have evolved alongside native plant communities and are often highly adapted to local climatic and environmental conditions, making them efficient pollinators of both wild plants and agricultural crops.

The ecological importance of native pollinators extends far beyond food production. Pollinator-mediated plant reproduction forms the basis of terrestrial food webs by supporting vegetation that provides food and shelter for numerous animal species. Healthy pollinator populations help maintain biodiversity, ecosystem productivity, and ecological resilience. In agricultural systems, pollinators increase crop yields, improve fruit quality, and enhance the nutritional diversity of food supplies [27-30]. Crops such as apples, almonds, berries, melons, cucumbers, and oilseeds rely heavily on pollinators for successful production. The economic value of pollination services is therefore immense and contributes significantly to global agricultural economies and rural livelihoods.

Despite their ecological and economic significance, native pollinator populations are declining worldwide. Habitat destruction, climate change, pesticide exposure, pollution, invasive species, and intensive agricultural practices have emerged as major threats to pollinator survival. The conversion of natural habitats into urban and agricultural landscapes has reduced the availability of nesting sites and floral resources necessary for pollinator reproduction and survival. Monoculture farming systems, while increasing short-term agricultural productivity, often fail to provide continuous and diverse food sources for pollinators throughout the year. As a result, many pollinator species experience nutritional deficiencies and population decline.

Pesticides represent another major factor contributing to pollinator losses. Chemical insecticides can directly kill pollinators or impair their ability to forage, navigate, communicate, and reproduce. Native pollinators are particularly vulnerable because they are often exposed to pesticides through contaminated nectar, pollen, soil, and water sources. In addition, climate

change is altering flowering seasons, temperature patterns, and rainfall cycles, disrupting the synchronized relationships between plants and pollinators. Such mismatches may reduce pollination efficiency and weaken ecological interactions within ecosystems.

The decline of native pollinators threatens biodiversity conservation and ecosystem resilience. Diverse pollinator communities are essential for maintaining stable ecological networks because different pollinator species respond differently to environmental changes and disturbances. A decline in pollinator diversity can reduce plant reproductive success, decrease genetic diversity, and destabilize ecosystems. Furthermore, the loss of pollinators may negatively affect wildlife species that depend on pollinated plants for food and habitat, thereby creating cascading ecological impacts throughout food chains.

2. Literature Review

Pollinators are among the most important biological agents sustaining terrestrial ecosystems and agricultural production systems. Their role in transferring pollen between flowers enables fertilization, seed production, and genetic diversity among flowering plants. However, growing scientific evidence suggests that pollinator populations are declining globally due to anthropogenic pressures such as habitat destruction, pesticide use, pollution, climate change, and the introduction of exotic species. These declines threaten ecosystem sustainability, biodiversity conservation, and global food security.

The relationship between pollinators and plants has been recognized since ancient times. Early observations by Theophrastus and Aristotle established preliminary links between insects and flowers. Later, Sprengel (1793) proposed that floral structures evolved in response to insect pollinators, while Darwin (1887) further demonstrated the evolutionary significance of cross-pollination. Contemporary ecological studies have expanded these theories by emphasizing the mutual dependence between pollinators and flowering plants in maintaining ecosystem stability.

Habitat alteration is considered one of the primary causes of pollinator decline. Agricultural intensification, urban expansion, monoculture farming, deforestation, and land fragmentation reduce the availability of floral resources and nesting sites for pollinators. According to Powell and Powell (1987), fragmentation significantly restricts pollinator movement, particularly among euglossine bees that avoid crossing large cleared areas. Such habitat barriers reduce genetic exchange among plant populations and limit pollination efficiency. Similarly, long-lived pollinators such as bumble bees suffer when agricultural practices remove shrubs, trees, and natural vegetation required for nesting and reproduction.

Monoculture agriculture also creates ecological imbalances by producing temporary floral abundance followed by prolonged food scarcity for pollinators. Crops such as sunflower, mustard, and alfalfa depend heavily on insect pollination, yet extensive monocultures often fail to support pollinator diversity year-round. Studies by Free (1964), Furgala et al. (1979), and Dhoble and Shinde (1982) demonstrated that insect pollination significantly improves seed yield and oil content in sunflower. Likewise, Kumar et al. (1994b) identified *Apis mellifera* and *Andrena ilderdia* as highly efficient pollinators of *Brassica campestris*. Despite these findings, modern agricultural systems continue to reduce natural habitats essential for sustaining pollinator populations outside crop flowering periods.

Pesticide exposure represents another major threat to pollinator survival. The increased use of chemical pesticides in intensive agriculture not only causes direct mortality but also results in sublethal effects that impair pollinator behavior, navigation, memory, and reproduction. Many developing countries continue to use pesticides indiscriminately, often exceeding recommended application levels. These chemicals contaminate pollen, nectar, and water sources,

thereby poisoning both managed and wild pollinator species. Research has shown that chronic pesticide exposure weakens bee colonies and contributes to long-term population declines.

Environmental pollution further compounds pollinator vulnerability. Heavy metals and airborne pollutants accumulate in floral tissues, pollen, and nectar through root uptake and atmospheric deposition. Pollinators such as honey bees subsequently ingest these contaminants during foraging activities. Studies indicate that pollen collected by bees can serve as bioindicators of environmental pollution because pollutant concentrations accumulate within bee bodies and hive products. Pollution stress negatively affects pollinator physiology, reproductive success, and overall ecosystem health.

Climate and environmental conditions also influence pollinator populations. Extreme rainfall, prolonged cold periods, and temperature fluctuations can disrupt pollinator emergence, flight activity, and breeding cycles. Soil-nesting bees are especially vulnerable to flooding and fungal infections caused by excessive soil moisture. According to ecological observations, severe weather conditions during flowering seasons often result in starvation and mortality among adult pollinators, thereby reducing pollination services in subsequent generations.

3. Methodology

Study Area

The study was conducted across selected agricultural, horticultural, and forest ecosystems in and around Chandigarh, Karnal, and Kinnaur. These locations were selected to examine pollinator communities under varying degrees of urbanization, habitat fragmentation, pesticide exposure, and agricultural intensification.

The Chandigarh tricity region represents a rapidly urbanizing agricultural ecosystem where monoculture farming, habitat modification, and pesticide use are common. Forest-associated studies were conducted near Madhuban, Karnal, while relatively undisturbed apple orchards in Kinnaur provided a comparative high-altitude ecosystem less affected by industrialization.

Selection of Plant Species

Six representative plant species belonging to agricultural, horticultural, and forest ecosystems were selected based on their ecological and economic importance as pollinator-dependent species.

- *Helianthus annuus* (Sunflower) – oilseed crop
- *Brassica campestris* var. sarson – oilseed and vegetable crop
- *Abelmoschus esculentus* (Okra) – vegetable crop
- *Malus domestica* var. royal delicious (Apple) – fruit crop
- *Litchi chinensis* (Litchi) – fruit crop
- *Eucalyptus tereticornis* – forest tree species

These crops differ in flowering duration, floral morphology, nectar production, and pollinator dependence, making them suitable for evaluating pollinator diversity and ecosystem interactions.

Survey of Pollinator Communities

Flower-visiting insects were sampled throughout the flowering period of each crop using standard sweep-net techniques. Collections were made every two hours between 0900 and 1700 hours during peak blooming stages. Opportunistic observations of pollinators outside scheduled intervals were also recorded.

Collected insects were killed using ethyl acetate and preserved following standard entomological procedures. Specimens were mounted, dried, and identified using taxonomic keys

and institutional reference collections from the Department of Zoology at Panjab University and Punjabi University.

Evaluation of Pollinator Decline Indicators

Pollinator decline was assessed using the following ecological indicators:

1. Species Diversity

The number of pollinator species associated with each crop was recorded to estimate diversity levels across habitats.

2. Relative Abundance

Five quadrats (1 m × 1 m) for field crops and five flowering branches for fruit crops were randomly selected. Pollinator visits were counted for five-minute intervals during three observation periods each day.

4. Results and Discussions

Pollinator Diversity and Foraging Behavior in Fruit Crops and Forest Flora

The investigation on apple (*Malus domestica*), litchi (*Litchi chinensis*), and *Eucalyptus tereticornis* revealed a diverse community of insect pollinators associated with fruit crops and forest flora. The results emphasized the ecological significance of insect-mediated pollination in sustaining fruit production, forest biodiversity, and ecosystem stability.

Apple Pollination

Apple orchards in Kinnaur supported a rich diversity of insect visitors during the flowering period. Ten insect species belonging to four insect orders were recorded visiting apple blossoms. The findings correspond closely with previous reports by Singh and Mishra (1986) and Mishra et al. (1976), who documented similar pollinator diversity in Himachal Pradesh apple orchards.

Among the recorded species, *Apis cerana* was the most abundant pollinator. The dominance of this native honey bee may be attributed to the absence of managed *Apis mellifera* colonies in the study area and the widespread traditional beekeeping practices maintained by local communities.




Figure 1: Performance ratings for insect pollinators' foraging duration on *Eucalyptus tereticornis* Sm.

Apis cerana demonstrated a high foraging rate (9.26 flowers/minute) and spent relatively less time per flower (5.53 seconds), indicating efficient pollination behavior. Similar observations were reported by Sharma (1961) and Kumar (1988), who identified *A. cerana* as the principal pollinator of apple orchards in Himalayan regions.

Andrena species carried the highest number of loose pollen grains (115,833 grains/insect), followed by *Apis cerana* (56,666 grains/insect). This finding suggests that solitary bees also contribute significantly to apple pollination, especially in high-altitude ecosystems.

The study confirmed that insect pollination is essential for apple production because most cultivars are self-incompatible and require cross-pollination for fruit set. Pollinator activity directly influenced fruit quality, seed development, and orchard productivity.

Litchi Pollination

Litchi flowers attracted a broad spectrum of insect visitors due to their strong nectar secretion and clustered floral arrangement. Honey bees, dipterans, and butterflies were the major floral visitors observed during the study.

Honey bees, particularly *Apis cerana* and *Apis mellifera*, were highly active on litchi inflorescences throughout the flowering period. Their frequent movement between flowers facilitated efficient cross-pollination, leading to improved fruit set.

The prolonged flowering period of litchi provided an important nectar source for pollinators. Similar observations were reported by Partap (1997), who identified litchi as an important honey plant supporting bee populations in subtropical regions.

The findings highlight the mutual relationship between pollinators and fruit crops. While pollinators obtain nectar and pollen resources, the crop benefits through enhanced fertilization and fruit production.

5. Conclusion

The investigation confirmed that insect pollinators are indispensable components of fruit crop production and forest ecosystem sustainability. Apple, litchi, and eucalyptus ecosystems supported a wide diversity of pollinating insects, with honey bees and solitary bees contributing most effectively to pollination. Among all pollinators studied, *Apis cerana* exhibited remarkable abundance and efficiency, particularly in apple orchards where cross-pollination is essential due to self-incompatibility among cultivars. Solitary bees such as *Andrena* species also demonstrated considerable pollination potential through their high pollen-carrying capacity. The study further revealed that flowering forest flora such as eucalyptus provides critical nectar resources that sustain pollinator populations and support apiculture activities. Pollinator foraging behavior, abundance, and diversity were strongly influenced by environmental conditions, habitat quality, and floral resource availability. The findings emphasized that the conservation of native pollinators and pollinator-friendly habitats is necessary not only for improving fruit yield and quality but also for preserving biodiversity and ecological balance. Increasing urbanization, habitat loss, climate variability, and excessive pesticide use pose serious threats to pollinator communities. Consequently, integrated conservation measures, ecological farming practices, and habitat restoration programs are urgently needed to sustain pollinator diversity and maintain healthy agro-ecosystems and forest ecosystems for future generations.

Acknowledgment: Not Applicable.

Funding Statement: The author(s) received no specific funding for this study.

Conflicts of Interest: The authors declare no conflicts of interest to report regarding the present study

References

- [1] D. P. Abrol, "Pollination studies in almond (*Prunus amygdalus* L.)," *Curr. Sci.*, vol. 57, pp. 397–398, 1988.
- [2] D. P. Abrol, "Insect pollination and crop production in Jammu and Kashmir," *Curr. Sci.*, vol. 65, no. 3, pp. 265–269, 1993.

-
- [3] A. R. Akhtar, "Studies in Indian Brassicae I: Sterility and selective pollen tube growth," *Indian J. Agric. Sci.*, vol. 2, pp. 280–292, 1932.
 - [4] P. Akwatanakul and M. Burgett, "Varroa jacobsoni: A perspective pest of honeybees in many parts of the world," *Bee World*, vol. 56, no. 3, pp. 119–121, 1975.
 - [5] A. H. Alex, "Pollination of some oilseed crops by honeybees," *Prog. Rep. Tex. Agric. Exp. Stn.*, pp. 1–5, 1960.
 - [6] G. Allen-Wardell, P. Bernhardt, R. Bitner, A. Burquez, S. Buchmann and J. Cane, "The potential consequences of pollinator declines on the conservation of biodiversity and stability of food crop yields," *Conserv. Biol.*, vol. 12, pp. 8–17, 1998.
 - [7] S. Appanah, "Pollination in Malaysian primary forests," *Malay. For.*, vol. 44, pp. 37–42, 1981.
 - [8] S. Appanah, "General flowering in the climax rain forests of Southeast Asia," *J. Trop. Ecol.*, vol. 1, pp. 225–240, 1985.
 - [9] S. Appanah, "Insect pollination and the diversity of dipterocarps," in *Proc. 3rd Roundtable Conf. Dipterocarps*, A. J. G. H. Kosterman, Ed., UNESCO, Jakarta, Java, 1987, pp. 277–291.
 - [10] S. Appanah and H. T. Chan, "Thrips: The pollinators of some dipterocarps," *Malay. For.*, vol. 44, pp. 234–252, 1981.
 - [11] D. R. Arya, R. C. Sihag and P. R. Yadav, "Diversity, abundance and foraging activity of insect pollinators of sunflower (*Helianthus annuus* L.) at Hisar (India)," *Indian Bee J.*, vol. 56, no. 3–4, pp. 172–178, 1994.
 - [12] D. R. Arya, R. C. Sihag and P. R. Yadav, "Role of insect pollination in seed yield of sunflower (*Helianthus annuus* L.)," *Indian Bee J.*, vol. 56, nos. 3–4, pp. 179–182, 1994.
 - [13] P. S. Ashton, T. J. Givnish and S. Appanah, "Staggered flowering in the Dipterocarpaceae: New insights into floral induction and the evolution of mast fruiting in the aseasonal tropics," *Am. Nat.*, vol. 132, pp. 44–66, 1988.
 - [14] A. S. Atwal and N. P. Goyal, "Introduction of *Apis mellifera* in Punjab plains," *Indian Bee J.*, vol. 35, pp. 1–9, 1973.
 - [15] C. C. Atwood, "Studies on the Apoidea of Western Nova Scotia with special reference to visitors to apple bloom," *Can. J. Res.*, vol. 9, pp. 443–457, 1933.
 - [16] S. D. Badiyala and R. Garg, "Studies on the pollination and fruit production by *Apis mellifera* L. in seven cultivars of litchi (*Litchi chinensis* Sonn.)," *Indian Bee J.*, vol. 52, pp. 28–30, 1990.
 - [17] H. G. Baker and P. D. Hurd, "Intrafloral ecology," *Annu. Rev. Entomol.*, vol. 13, pp. 385–414, 1968.
 - [18] J. Banaszak, "Strategy for conservation of wild bees in an agricultural landscape," *Agric., Ecosyst. Environ.*, vol. 40, pp. 179–192, 1992.
 - [19] S. W. T. Batra, "Crop pollination and the flower relationships of the wild bees of Ludhiana, India (Hymenoptera: Apoidea)," *J. Kans. Entomol. Soc.*, vol. 40, no. 2, pp. 164–177, 1967.
 - [20] E. I. Belozera, "Bees increase the seed crop from winter rape," *Pchelovodstvo*, vol. 37, no. 9, pp. 38–40, 1960.
 - [21] P. Benedek, "Structure of flower visiting insect populations and effect of environmental conditions on honeybee density in blooming winter rape fields and their bearing on pest control," *Novénytermelés*, vol. 25, pp. 65–74, 1976.
 - [22] P. Benedek, "Economic importance of honey pollination of crops at the national level in Hungary," in *Proc. 29th Int. Congr. Apiculture—Apimondia*, Bucharest, Romania, 1985, pp. 286–289.
 - [23] P. Benedek and J. Prenner, "Effect of temperature on the behaviour and pollinating efficiency of honey bees on winter rape flowers," *Z. Angew. Entomol.*, vol. 71, no. 2, pp. 120–124, 1972.
-

-
- [24] O. P. Bhalla, A. K. Verma, and H. S. Dhaliwal, "Insect visitors of mustard bloom (*Brassica campestris* var. *sarson*), their number and foraging behaviour under mid-hill conditions," *J. Entomol. Res.*, vol. 7, no. 1, pp. 16–17, 1983.
- [25] O. P. Bhalla, A. K. Verma, and H. S. Dhaliwal, "Foraging activity of insect pollinators visiting stone fruits," *J. Entomol. Res.*, vol. 7, no. 2, pp. 91–94, 1983.
- [26] P N S S V Charishma, S. Rupa Lakshmi, and M. Vijaya Durga, "Making Crop Recommendations using Machine Learning Techniques," *JCAI*, vol. 2, no. 2, pp. 1–12, 2024.
- [27] G. E. Bohart, "Insect pollination of forage legumes," *Bee World*, vol. 41, pp. 57–64, 85–97, 1960.
- [28] P. Brundavani, "A Principal Component Analysis Algorithm for Seed Enterprise Financial Performance and Scientific and Technological Innovation," *JCAI*, vol. 2, no. 2, pp. 49–62, 2024.
- [29] R. M. Borges, "Leviathan, natural selection and ethics," *Curr. Sci.*, vol. 74, pp. 750–758, 1999.
- [30] R. M. D. Boyle and B. J. R. Philogene, "The native pollinators of an apple orchard, variations and significance," *J. Hortic. Sci.*, vol. 58, no. 3, pp. 355–363, 1983.